

Product datasheet for TA323925S

uPA (PLAU) Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IHC
Recommended Dilution:	IHC: 50-200 Positive control: Human liver cancer Predicted cell location: Cytoplasm
Reactivity:	Human
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Fusion protein corresponding to a region derived from 179-417 amino acids of human plasminogen activator, urokinase
Formulation:	PBS pH7.3, 0.05% NaN3, 50% glycerol
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	plasminogen activator, urokinase
Database Link:	<u>NP_002649</u> <u>Entrez Gene 5328 Human</u> <u>P00749</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Section 2012 CRIGENE uPA (PLAU) Rabbit Polyclonal Antibody – TA323925S

Background: This gene encodes a serine protease involved in degradation of the extracellular matrix and possibly tumor cell migration and proliferation. A specific polymorphism in this gene may be associated with late-onset Alzheimer's disease and also with decreased affinity for fibrin-binding. This protein converts plasminogen to plasmin by specific cleavage of an Arg-Val bond in plasminogen. Plasmin in turn cleaves this protein at a Lys-Ile bond to form a two-chain derivative in which a single disulfide bond connects the amino-terminal A-chain to the catalytically active; carboxy-terminal B-chain. This two-chain derivative is also called HMW-uPA (high molecular weight uPA). HMW-uPA can be further processed into LMW-uPA (low molecular weight uPA) by cleavage of chain A into a short chain A (A1) and an amino-terminal fragment. LMW-uPA is proteolytically active but does not bind to the uPA receptor. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Synonyms:ATF; BDPLT5; QPD; u-PA; UPA; URKProtein Families:Druggable Genome, ES Cell Differentiation/IPS, ProteaseProtein Pathways:Complement and coagulation cascades

Product images:



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA323925] (PLAU Antibody) at dilution 1/70 (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA323925] (PLAU Antibody) at dilution 1/70, treated with fusion protein. (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA323925] (PLAU Antibody) at dilution 1/70 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA323925] (PLAU Antibody) at dilution 1/70, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US