

Product datasheet for TA323961

CDK5 Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type: | Primary Antibodies |
|-------------------------|---|
| Applications: | IHC, WB |
| Recommended Dilution: | WB: 500-2000 WB positive control: Jurkat and NIH/3T3 cells IHC: 25-100 Positive control: Human colon cancer Predicted cell location: Cytoplasm or Nucleus |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| lsotype: | lgG |
| Clonality: | Polyclonal |
| Immunogen: | Full length fusion protein |
| Formulation: | PBS pH7.3, 0.05% NaN3, 50% glycerol |
| Concentration: | lot specific |
| Purification: | Antigen affinity purification |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Predicted Protein Size: | 33 kDa |
| Gene Name: | cyclin-dependent kinase 5 |
| Database Link: | <u>NP_001157882</u> Entrez Gene 12568 MouseEntrez Gene 140908 RatEntrez Gene 1020 Human Q00535 |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


CDK5 Rabbit Polyclonal Antibody – TA323961

Background:Cell division protein kinase 5?is an?enzyme?that in humans is encoded by the?CDK5?gene.?
The protein encoded by this gene is part of the?cyclin-dependent kinase?family. CDK5 is
required for proper development of the brain and to be activated; CDK5 must associate with
CDK5R1 or CDK5R2. Cdk5 is involved in the processes of neuronal maturation and migration;
phosphorylating the key intracellular adaptor of the reelin signaling chain. Experiments
performed on mice lacking p35; a necessary activator of cdk5 in early brain development;
showed that the normal layering of neurons was reversed in the cortex. This disrupted
lamination again implicated cdk5 in neuronal migration and plasticity. Cdk5 is also involved in
the regulation of synaptic vesicle exocytosis via phosphorylation of munc-18.

Synonyms: PSSALRE


Protein Families:Druggable Genome, Protein KinaseProtein Pathways:Alzheimer's disease, Axon guidance

Product images:



Gel: 10%SDS-PAGE Lysate: 40 µg Lane 1-2: Jurkat cells NIH/3T3 cells Primary antibody: TA323961 (CDK5 Antibody) at dilution 1/300 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 1 minute

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US




Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA323961 (CDK5 Antibody) at dilution 1/30 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA323961 (CDK5 Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human brain tissue using TA323961 (CDK5 Antibody) at dilution 1/30 (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human brain tissue using TA323961 (CDK5 Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US