

# Product datasheet for TA323962S

## **CDK5 Rabbit Polyclonal Antibody**

### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 200-1000 WB positive control: Human fetal brain and colon cancer tissue IHC: 10-50 Positive control: Human liver cancer Predicted cell location: Cytoplasm
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Full length fusion protein
Formulation:	PBS pH7.3, 0.05% NaN3, 50% glycerol
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	33 kDa
Gene Name:	cyclin-dependent kinase 5
Database Link:	<u>NP_001157882</u> <u>Entrez Gene 12568 MouseEntrez Gene 140908 RatEntrez Gene 1020 Human</u> <u>Q00535</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **CDK5** Rabbit Polyclonal Antibody – TA323962S

Background:Cell division protein kinase 5?is an?enzyme?that in humans is encoded by the?CDK5?gene.?<br/>The protein encoded by this gene is part of the?cyclin-dependent kinase?family. CDK5 is<br/>required for proper development of the brain and to be activated; CDK5 must associate with<br/>CDK5R1 or CDK5R2. Cdk5 is involved in the processes of neuronal maturation and migration;<br/>phosphorylating the key intracellular adaptor of the reelin signaling chain. Experiments<br/>performed on mice lacking p35; a necessary activator of cdk5 in early brain development;<br/>showed that the normal layering of neurons was reversed in the cortex. This disrupted<br/>lamination again implicated cdk5 in neuronal migration and plasticity. Cdk5 is also involved in<br/>the regulation of synaptic vesicle exocytosis via phosphorylation of munc-18.


Synonyms:	PSSALRE
Protein Families:	Druggable Genome, Protein Kinase
Protein Pathways:	Alzheimer's disease, Axon guidance

#### **Product images:**



Gel: 10+12%SDS-PAGE Lysate: 30 µg Lane 1-2: Human fetal brain tissue Human colon cancer tissue Primary antibody: [TA323962] (CDK5 Antibody) at dilution 1/150 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 5 minutes

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA323962] (CDK5 Antibody) at dilution 1/10 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA323962] (CDK5 Antibody) at dilution 1/10, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US