

Product datasheet for TA324020S

VEGFA Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: 293T cell lysate

IHC: 25-50

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 130-143 amino acids of human

vascular endothelial growth factor A

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: vascular endothelial growth factor A

Database Link: NP 001165094

Entrez Gene 7422 Human

P15692



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

This gene is a member of the PDGF/VEGF growth factor family and encodes a protein that is often found as a disulfide linked homodimer. This protein is a glycosylated mitogen that specifically acts on endothelial cells and has various effects; including mediating increased vascular permeability; inducing angiogenesis; vasculogenesis and endothelial cell growth; promoting cell migration; and inhibiting apoptosis. Elevated levels of this protein is linked to POEMS syndrome; also known as Crow-Fukase syndrome. Mutations in this gene have been associated with proliferative and nonproliferative diabetic retinopathy. Alternatively spliced transcript variants; encoding either freely secreted or cell-associated isoforms; have been characterized. There is also evidence for the use of non-AUG (CUG) translation initiation sites upstream of; and in-frame with the first AUG; leading to additional isoforms.

Synonyms: MGC70609; MVCD1; VEGF; VPF


Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Bladder cancer, Cytokine-cytokine receptor interaction, Focal adhesion, mTOR signaling

pathway, Pancreatic cancer, Pathways in cancer, Renal cell carcinoma, VEGF signaling

pathway

Product images:



Gel: 12%SDS-PAGE Lysate: 40 µg Lane: 293T cell lysate

Primary antibody: [TA324020] (VEGFA Antibody)


at dilution 1/400

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 3 minutes





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA324020] (VEGFA Antibody) at dilution 1/25 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA324020] (VEGFA Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)