

Product datasheet for TA324064S

EGF Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human prostate tissue lysate

IHC: 100-300

Positive control: Human cervical cancer

Predicted cell location: Cytoplasm and Cell membrane

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to a region derived from 926-1026 amino acids of human

epidermal growth factor

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 134 kDa

Gene Name: epidermal growth factor

Database Link: NP 001171601

Entrez Gene 1950 Human

P01133



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

EGF Rabbit Polyclonal Antibody - TA324064S

Background: This gene encodes a member of the epidermal growth factor superfamily. The encoded

protein is synthesized as a large precursor molecule that is proteolytically cleaved to generate the 53-amino acid epidermal growth factor peptide. This protein acts a potent mitogenic factor that plays an important role in the growth; proliferation and differentiation of numerous cell types. This protein acts by binding the high affinity cell surface receptor; epidermal growth factor receptor. Defects in this gene are the cause of hypomagnesemia type 4. Dysregulation of this gene has been associated with the growth and progression of certain cancers. Alternate splicing results in multiple transcript variants

Synonyms: HOMG4; URG

Protein Families: Adult stem cells, Druggable Genome, Embryonic stem cells, ES Cell Differentiation/IPS,


Induced pluripotent stem cells, Transmembrane

Protein Pathways: Bladder cancer, Cytokine-cytokine receptor interaction, Endocytosis, Endometrial cancer, ErbB

signaling pathway, Focal adhesion, Gap junction, Glioma, MAPK signaling pathway, Melanoma, Non-small cell lung cancer, Pancreatic cancer, Pathways in cancer, Prostate

cancer, Regulation of actin cytoskeleton

Product images:



Gel: 6%SDS-PAGE Lysate: 40 μg

Lane: Human prostate tissue lysate

Primary antibody: [TA324064] (EGF Antibody) at


dilution 1/700

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 30 seconds





Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA324064] (EGF Antibody) at dilution 1/140 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA324064] (EGF Antibody) at dilution 1/140, treated with fusion protein. (Original magnification: ×200)