

## **Product datasheet for TA324083S**

## Product datasireet for TA3240653

## Product data:

**Product Type:** Primary Antibodies

**SIRT2 Rabbit Polyclonal Antibody** 

Applications: WB

Recommended Dilution: WB: 200-1000

WB positive control: Mouse heart tissue

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein corresponding to a region derived from 65-340 amino acids of human sirtuin

2

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 43 kDa **Gene Name:** sirtuin 2

Database Link: NP 001180215

Entrez Gene 64383 MouseEntrez Gene 361532 RatEntrez Gene 22933 Human

Q8IXJ6

**Background:** This gene encodes a member of the sirtuin family of proteins; homologs to the yeast Sir2

protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however; yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Several transcript variants are resulted from

alternative splicing of this gene.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**Synonyms:** SIR2; SIR2L; SIR2L2

**Protein Families:** Druggable Genome, Transcription Factors

## **Product images:**



Gel: 10%SDS-PAGE Lysate: 40 μg

Lane: Mouse heart tissue

Primary antibody: [TA324083] (SIRT2 Antibody) at

dilution 1/350

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 30 seconds