

Product datasheet for TA324329S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Lactoferrin (LTF) Mouse Monoclonal Antibody [Clone ID: 119CT80.1.1]

Product data:

Product Type: Primary Antibodies

Clone Name: 119CT80.1.1
Applications: IF, IHC, WB

Recommended Dilution: WB: 1:1000, IHC: 1:50~100, IF: 1:10~50

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: This LTF Monoclonal antibody is generated from mouse immunized with LTF recombinant

protein.

Formulation: Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is purified through a protein G column, followed by dialysis against PBS.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 78182 Da

Gene Name: lactotransferrin

Database Link: NP 002334

Entrez Gene 4057 Human

P02788


Synonyms: GIG12; HEL110; HLF2; LF

Protein Families: Druggable Genome, Protease, Secreted Protein





Product images:





Western blot analysis of lysate from K562 cell line, using LTF Antibody (Cat. #[TA324329]). [TA324329] was diluted at 1:1000. A goat antimouse IgG H&L (HRP) at 1:3000 dilution was used as the secondary antibody. Lysate at 35ug.



Formalin-fixed and paraffin-embedded human prostate carcinoma with LTF Monoclonal Antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.







Immunofluorescence analysis of LTF Monoclonal Antibody with paraffin-embedded human prostate carcinoma tissue. 0.05 mg/ml primary antibody was followed by PE-conjugated goat anti-mouse lgG (whole molecule). PE emits orange fluorescence.

Confocal immunofluorescent analysis of LTF Antibody (Cat#[TA324329]) with HepG2 cell followed by Alexa Fluor® 488-conjugated goat anti-mouse IgG (green). DAPI was used to stain the cell nuclear (blue).