

Product datasheet for TA324354

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

PINK1 Mouse Monoclonal Antibody [Clone ID: 38CT20.8.5]

Product data:

Product Type: Primary Antibodies

Clone Name: 38CT20.8.5
Applications: IHC, WB

Recommended Dilution: WB: 1:500~2000, IHC: 1:10~50

Reactivity: Human, Mouse

Host: Mouse

Clonality: Monoclonal

Immunogen: Recombinant PINK1 protein was used to produced this monoclonal antibody.

Formulation: Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.

Concentration: lot specific

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 62769 Da

Gene Name: PTEN induced putative kinase 1

Database Link: NP 115785

Entrez Gene 68943 MouseEntrez Gene 65018 Human

Q9BXM7


Synonyms: BRPK; PARK6

Protein Families: Druggable Genome, Protein Kinase


Protein Pathways: Parkinson's disease



Product images:





Western blot analysis of anti-PINK1 Monoclonal Antibody (TA324354) in mouse brain tissue lysates. PINK1 (arrow) was detected using the ascites Mab. (dilution 1:500)



Western blot analysis of PINK (arrow) using mouse monoclonal PINK antibody (Ascites). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the PINK gene (Lane 2) (Origene Technologies) (1:2000)







Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with PINK1 Monoclonal Antibody (Cat.#TA324354), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Formalin-fixed and paraffin-embedded human Brain Cortex tissue reacted with PINK1 Monoclonal Antibody (Cat.#TA324354), which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.