

Product datasheet for TA324367

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Folate Binding Protein (FOLR1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: FC, IHC, WB

Recommended Dilution: WB: 1:1000, IHC: 1:10~50, FC: 1:10~50

Reactivity: Human Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: This FOLR1 antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 33-68 amino acids from the N-terminal region of human FOLR1.

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is purified through a protein A column, followed by peptide affinity purification.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 29819 Da

Gene Name: folate receptor 1 (adult)

Database Link: NP 000793

Entrez Gene 2348 Human

P15328


Synonyms: FBP; FOLR

Protein Families: Druggable Genome, Secreted Protein, Transmembrane






Product images:




Western blot analysis of lysate from Hela cell line, using FOLR1 Antibody (N-term) (Cat. #TA324367). TA324367 was diluted at 1:1000. A goat antirabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.



FOLR1 antibody (N-term) (Cat. #TA324367) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the FOLR1 antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.





FOLR1 Antibody (N-term) (Cat. #TA324367) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.