

Product datasheet for TA324475

PCYT1A Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: IHC: 1:10~50, WB: 1:1000, IF: 1:10~50

Reactivity: Human (Predicted: Mouse, Rat, Hamster)

Host: Rabbit

Isotype: lgG

Clonality: Polyclonal

Immunogen: This PCYT1A antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 27-54 amino acids from the N-terminal region of human PCYT1A.

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is purified through a protein A column, followed by peptide affinity purification.

Conjugation: Unconjugated

Store at -20°C as received. Storage:

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 41731 Da

Gene Name: phosphate cytidylyltransferase 1, choline, alpha

Database Link: NP 005008

Entrez Gene 13026 MouseEntrez Gene 140544 RatEntrez Gene 5130 Human

P49585

Synonyms: CCTA; CT; CTA; CTPCT; PCYT1; SMDCRD

Protein Pathways: Glycerophospholipid metabolism, Metabolic pathways




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:




PCYT1A Antibody (N-term) (Cat. #TA324475) western blot analysis in K562, MDA-MB453 cell line lysates (35ug/lane). This demonstrates the PCYT1A antibody detected the PCYT1A protein (arrow).



PCYT1A Antibody (N-term) (Cat. #TA324475)immunohistochemistry analysis in formalin fixed and paraffin embedded human colon carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of PCYT1A Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.





Confocal immunofluorescent analysis of PCYT1A Antibody (N-term) (Cat#TA324475) with WiDr cell followed by Alexa Fluor 488-conjugated goat antirabbit IgG (green). DAPI was used to stain the cell nuclear (blue).