

Product datasheet for TA324505

Sumo 3 (SUMO3) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: WB: 1:1000, IF: 1:10~50, IHC: 1:50~100

Reactivity: Human, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: This Pan SUMO antibody recognizes all 3 SUMO isoforms, including human SUMO1, SUMO2

and SUMO3. This antibody is generated from rabbits immunized with a recombinant protein

encoding full length human SUMO3.

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by

dialysis against PBS.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 11637 Da

Gene Name: small ubiquitin-like modifier 3

Database Link: NP 008867

Entrez Gene 499417 RatEntrez Gene 6612 Human

P55854

Synonyms: SMT3A; Smt3B; SMT3H1; SUMO-3

Protein Families: Druggable Genome




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:




SUMO3 Antibody (Cat. #TA324505) western blot analysis in 293 cell line and rat liver tissue lysates (35ug/lane). This demonstrates the SUMO3 antibody detected the SUMO3 protein (arrow).



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.





IF image of Hela cell stained with Pan SUMO Antibody (Cat#TA324505). Hela cells were incubated with Pan SUMO primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400). Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7 units/ml). Nuclei were counterstained with DAPI (blue). Pan SUMO immunoreactivity is localized to Nucleus significantly.