

Product datasheet for TA324512

NFATC4 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, WB

Recommended Dilution: WB: 1:1000, IF: 1:10~50

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: This NFATC4 antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 744-773 amino acids from the C-terminal region of human NFATC4.

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is purified through a protein A column, followed by peptide affinity purification.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 95449 Da

Gene Name: nuclear factor of activated T-cells 4

Database Link: NP 001129494

Entrez Gene 4776 Human

Q14934

Synonyms: NF-AT3; NF-ATC4; NFAT3

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Axon guidance, B cell receptor signaling pathway, MAPK signaling pathway, Natural killer cell

mediated cytotoxicity, T cell receptor signaling pathway, VEGF signaling pathway, Wnt

signaling pathway




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



NFATC4 Antibody (C-term) (Cat. #TA324512) western blot analysis in human placenta tissue lysates (35ug/lane). This demonstrates the NFATC4 antibody detected the NFATC4 protein (arrow).



IF image of U251 cell stained with NFATC4 Antibody (C-term) (Cat#TA324512).U251 cells were incubated with NFATC4 primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400).Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7 units/ml). Nuclei were counterstained with DAPI (blue). NFATC4 immunoreactivity is localized to Cytoplasm significantly.