

Product datasheet for TA324729

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Vimentin (VIM) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: FC, IF, IHC, WB

Recommended Dilution: IF: 1:100, WB: 1:1000, IHC: 1:10~50, FC: 1:10~50

Reactivity: Human (Predicted: Mouse, Rat, Bovine, Chicken, Hamster, Pig, Monkey, Xenopus)

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: This Vimentin antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 430-457 amino acids from the C-terminal region of human Vimentin.

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by

dialysis against PBS.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 53652 Da

Gene Name: vimentin

Database Link: NP 003371

Entrez Gene 22352 MouseEntrez Gene 81818 RatEntrez Gene 705289 MonkeyEntrez Gene

<u>7431 Human</u>

P08670


Synonyms: CTRCT30; HEL113

Protein Families: ES Cell Differentiation/IPS






Product images:





Western blot analysis of Vimentin antibody (Cterm) (Cat.#TA324729) in NCI-H460 cell line lysates (35ug/lane). Vimentin (arrow) was detected using the purified Pab.



Vimentin Antibody (C-term) (Cat. #TA324729)immunohistochemistry analysis in formalin fixed and paraffin embedded human colon tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of Vimentin Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.







IF image of SY5Y cells stained with Vimentin (Cterm) antibody. SY5Y cells were incubated with TA324729 Vimentin (C-term) primary antibody (1:100, 2 h at RT). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:1000, 1h). Nuclei were counterstained with Hoechst 33342 (blue). Note the highly specific localization of the Vimentin immunosignal to the cytoskeleton, supported by Human Protein Atlas Data (http://www.proteinatlas.org/ENSG00000026025).

Vimentin Antibody (C-term) (Cat. #TA324729) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.