

## **Product datasheet for TA324746**

## Froduct datasireet for TA32474

## **FEN1 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IF, WB

Recommended Dilution: IF: 1:100, WB: 1:1000

Reactivity: Human (Predicted: Bovine)

**Host:** Rabbit

**Isotype:** IgG

Clonality: Polyclonal

Immunogen: This FEN1 antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 243-272 amino acids from the Central region of human FEN1.

**Formulation:** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

**Concentration:** lot specific

**Purification:** This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by

dialysis against PBS.

**Conjugation:** Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 42593 Da

**Gene Name:** flap structure-specific endonuclease 1

Database Link: NP 004102

Entrez Gene 2237 Human

P39748

**Synonyms:** FEN-1; MF1; RAD2

**Protein Families:** Druggable Genome, Stem cell - Pluripotency

**Protein Pathways:** Base excision repair, DNA replication, Non-homologous end-joining




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**




Western blot analysis of anti-FEN1 Antibody (Center) (Cat.#TA324746) in HepG2 cell line lysates (35ug/lane). FEN1 (arrow) was detected using the purified Pab.



Western blot analysis of FEN1 (arrow) using rabbit polyclonal FEN1 Antibody (Center) (Cat.#TA324746). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the FEN1 gene.





Immunofluorescent analysis of U251 cells, using FEN1 Antibody (Center) (Cat. #TA324746). TA324746 was diluted at 1:100 dilution. Alexa Fluor 488-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody (green). DAPI was used to stain the cell nuclear (blue).