

Product datasheet for TA324776S

p53 (TP53) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: Dot, IHC, WB

Recommended Dilution: DB: 1:500, WB: 1:1000, IHC: 1:50~100

Reactivity: Human, Mouse (Predicted: Pig, Monkey, Rabbit)

Modifications: Phospho-specific

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: This p53 Antibody is generated from rabbits immunized with a KLH conjugated synthetic

phosphopeptide corresponding to amino acid residues surrounding T18 of human p53.

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is purified through a protein A column, followed by peptide affinity purification.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 43653 Da

Gene Name: tumor protein p53

Database Link: NP 000537

Entrez Gene 22059 MouseEntrez Gene 716170 MonkeyEntrez Gene 7157 Human

P04637

Synonyms: BCC7; LFS1; P53; TRP53

Protein Families: Druggable Genome, Stem cell - Pluripotency, Transcription Factors



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Protein Pathways:

Amyotrophic lateral sclerosis (ALS), Apoptosis, Basal cell carcinoma, Bladder cancer, Cell cycle, Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, Glioma, Huntington's disease, MAPK signaling pathway, Melanoma, Neurotrophin signaling pathway, Non-small cell lung cancer, p53 signaling pathway, Pancreatic cancer, Pathways in cancer, Prostate cancer, Small cell lung cancer, Thyroid cancer, Wnt signaling pathway

Product images:




Western blot analysis of anti-Phospho-p53-T18 Pab (cat# [TA324776]) in, from left to right, A2058, Ramos, mouse lung, mouse testis, and HL60 cell line lysates. Phospho-p53-T18 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.





Dot blot analysis of anti-hp53-T18 Phosphospecific Pab (Cat. #[TA324776]) on nitrocellulose membrane. 50ng of nonphospho-peptide or phospho-peptide were adsorbed on their respective dots. Antibody working concentration was 0.5ug per ml.