

Product datasheet for TA324833

Kallikrein 6 (KLK6) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: FC, IHC, WB

Recommended Dilution: WB: 1:1000, IHC: 1:10~50, FC: 1:10~50

Reactivity: Rabbit Host: Isotype: **IgG**

Clonality: Polyclonal

Immunogen: This Kallikrein 6 antibody is generated from rabbits immunized with a KLH conjugated

synthetic peptide between 126-156 amino acids from the Central region of human Kallikrein

6.

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by

dialysis against PBS.

Conjugation: Unconjugated

Store at -20°C as received. Storage:

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 26856 Da

Gene Name: kallikrein related peptidase 6

Database Link: NP 001012982

Entrez Gene 5653 Human

Q92876

Bssp; hK6; Klk7; PRSS9; PRSS18; SP59 Synonyms:

Protein Families: Druggable Genome, Protease, Secreted Protein




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:




Western blot analysis of KLK6 (arrow) using rabbit polyclonal KLK6 Antibody (L141) (Cat. #TA324833). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the KLK6 gene.



Kallikrein 6 (KLK6) Antibody (Center) (Cat. #TA324833)immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of Kallikrein 6 (KLK6) Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.





Kallikrein 6 (KLK6) Antibody (Center) Antibody (Cat. #TA324833) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goatanti-rabbit secondary antibodies were used for the analysis.