

Product datasheet for TA324844

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Endothelin A Receptor (EDNRA) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: FC, IF, WB

Recommended Dilution: WB: 1:1000, IF: 1:10~50, FC: 1:10~50

Reactivity: Human (Predicted: Mouse, Rat)

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: This EDNRA antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 125-154 amino acids from the Central region of human EDNRA.

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by

dialysis against PBS.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 48722 Da

Gene Name: endothelin receptor type A

Database Link: NP 001159527

Entrez Gene 13617 MouseEntrez Gene 24326 RatEntrez Gene 1909 Human

P25101

Synonyms: ET-A; ETA; ETA-R; ETAR; ETRA; hET-AR

Protein Families: Druggable Genome, GPCR, Transmembrane


Protein Pathways: Calcium signaling pathway, Neuroactive ligand-receptor interaction, Vascular smooth muscle

contraction






Product images:




Western blot analysis of EDNRA antibody (Center) (Cat.# TA324844) in CEM cell line lysates (35ug/lane). EDNRA (arrow) was detected using the purified Pab.



Confocal immunofluorescent analysis of EDNRA Antibody (Center) (Cat#TA324844) with HepG2 cell followed by Alexa Fluoræ´?488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).





Flow cytometric analysis of CEM cells using EDNRA Antibody (Center) (green histogram) compared to a negative control cell (blue histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.