

Product datasheet for TA324864

Product data:

Product Type: Primary Antibodies

ADH1B Rabbit Polyclonal Antibody

Applications: FC, IF, IHC, WB

Recommended Dilution: WB: 1:1000, IHC: 1:50~100, FC: 1:10~50, IF: 1:10~50

Reactivity: Human (Predicted: Mouse, Rat, Chicken)

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: This ADH1B antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 209-237 amino acids from the Central region of human ADH1B.

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by

dialysis against PBS.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 39855 Da

Gene Name: alcohol dehydrogenase 1B (class I), beta polypeptide

Database Link: NP 000659

Entrez Gene 125 Human

P00325

Synonyms: ADH2; HEL-S-117

Protein Families: Druggable Genome

Protein Pathways: Drug metabolism - cytochrome P450, Fatty acid metabolism, Glycolysis / Gluconeogenesis,

Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Retinol metabolism,

Tyrosine metabolism




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Product images:



Western blot analysis of ADH1B Antibody (Center) (Cat. #TA324864) in HepG2 cell line lysates (35ug/lane). ADH1B (arrow) was detected using the purified Pab.




Formalin-fixed and paraffin-embedded human hepatocarcinoma reacted with ADH1B Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Confocal immunofluorescent analysis of ADH1B Antibody (Center) (Cat#TA324864) with HepG2 cell followed by Alexa Fluor® 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).





ADH1B Antibody (Center) (Cat.#TA324864) flow cytometry analysis of HepG2 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.