

Product datasheet for TA324880S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Ornithine Carbamoyltransferase (OTC) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: FC, IHC, WB

Recommended Dilution: WB: 1:1000, IHC: 1:10~50, FC: 1:10~50

Reactivity: Human (Predicted: Mouse, Rat, Bovine, Pig)

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: This OTC antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 71-98 amino acids from the Central region of human OTC.

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is purified through a protein A column, followed by peptide affinity purification.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 39935 Da

Gene Name: ornithine carbamoyltransferase

Database Link: NP 000522

Entrez Gene 18416 MouseEntrez Gene 25611 RatEntrez Gene 5009 Human

P00480

Synonyms: OCTD


Protein Families: Druggable Genome

Protein Pathways: Arginine and proline metabolism, Metabolic pathways







Product images:



OTC Antibody (Center) (Cat.# [TA324880]) western blot analysis in SW620 cell line and mouse liverà rat liver lysates (35ug/lane). This demonstrates the OTC antibody detected the OTC protein (arrow).



Formalin-fixed and paraffin-embedded human hepatoarcinoma with OTC Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of ATDC5 cells using OTC Antibody (Center) (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.