

Product datasheet for TA324948S

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MCSF Receptor (CSF1R) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: FC, IF, WB

Recommended Dilution: WB: 1:1000, IF: 1:10~50, FC: 1:10~50

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: This CSF1R antibody is generated from rabbits immunized with CSF1R recombinant protein.

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is purified through a protein A column, followed by peptide affinity purification.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 107984 Da

Gene Name: colony stimulating factor 1 receptor

Database Link: NP 005202

Entrez Gene 1436 Human

P07333

Synonyms: C-FMS; CD115; CSF-1R; CSFR; FIM2; FMS; HDLS; M-CSF-R

Protein Families: Druggable Genome, Protein Kinase, Transmembrane


Protein Pathways: Cytokine-cytokine receptor interaction, Endocytosis, Hematopoietic cell lineage, Pathways in

cancer






Product images:




Western blot analysis of lysate from U-937 cell line, using CSF1R Antibody (Cat. #[TA324948]). [TA324948] was diluted at 1:1000. A goat antirabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.



Confocal immunofluorescent analysis of CSF1R Antibody (Cat#[TA324948]) with HepG2 cell followed by Alexa Fluor 488-conjugated goat antirabbit IgG (green). DAPI was used to stain the cell nuclear (blue).





CSF1R Antibody (Cat. #[TA324948]) flow cytometric analysis of k562 cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.