

Product datasheet for TA324962S


FLT3 Rabbit Polyclonal Antibody

Product data:


Product Type:	Primary Antibodies
Applications:	FC, IHC, WB
Recommended Dilution:	FC: 1:10~50, WB: 1:1000, IHC: 1:50~100
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	This FLT3 (CD135) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 33-64 amino acids from the N-terminal region of human FLT3 (CD135).
Formulation:	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Concentration:	lot specific
Purification:	This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	112903 Da
Gene Name:	fms related tyrosine kinase 3
Database Link:	NP_004110 Entrez Gene 2322 Human P36888
Synonyms:	CD135; FLK-2; FLK2; STK1
Protein Families:	Druggable Genome, ES Cell Differentiation/IPS, Protein Kinase, Transmembrane
Protein Pathways:	Acute myeloid leukemia, Cytokine-cytokine receptor interaction, Hematopoietic cell lineage, Pathways in cancer


[View online »](#)


Product images:



Western blot analysis of anti-FLT3 Pab (Cat. # [TA324962]) in HL-60 cell lysate. FLT3 (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



FLT3 (CD135) Antibody (N-term) (Cat. # [TA324962]) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.