

## **Product datasheet for TA325044**

## **VAP1 (AOC3) Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** FC, IHC, WB

**Recommended Dilution:** WB: 1:1000, IHC: 1:10~50, FC: 1:10~50

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

**Immunogen:** This AOC3 antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 613-640 amino acids from the Central region of human AOC3.

**Formulation:** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

**Concentration:** lot specific

**Purification:** This antibody is purified through a protein A column, followed by peptide affinity purification.

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 84622 Da

**Gene Name:** amine oxidase, copper containing 3

Database Link: NP 003725

Entrez Gene 8639 Human

Q16853

Synonyms: HPAO; SSAO; VAP-1; VAP1

**Protein Families:** Transmembrane

**Protein Pathways:** beta-Alanine metabolism, Glycine, serine and threonine metabolism, Metabolic pathways,

Phenylalanine metabolism, Tyrosine metabolism




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**




Western blot analysis of AOC3 Antibody (Center) (Cat. #TA325044) in CEM cell line lysates (35ug/lane).AOC3 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human colon carcinoma with AOC3 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.





Flow cytometric analysis of CEM cells using AOC3 Antibody (Center) (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.