

Product datasheet for TA325117S

PAFAH (PLA2G7) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: FC, IHC, WB

Recommended Dilution: WB: 1:1000, IHC: 1:10~50, FC: 1:10~50

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: This PLA2G7 antibody is generated from rabbits immunized with a KLH conjugated synthetic

peptide between 200-228 amino acids from the Central region of human PLA2G7.

Formulation: Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Concentration: lot specific

Purification: This antibody is purified through a protein A column, followed by peptide affinity purification.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 50077 Da

Gene Name: phospholipase A2 group VII

Database Link: NP 005075

Entrez Gene 7941 Human

Q13093

Synonyms: LDL-PLA2; LP-PLA2; PAFAD; PAFAH

Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Ether lipid metabolism, Metabolic pathways




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:




Western blot analysis of PLA2G7 Antibody (Center) (Cat. #[TA325117]) in HL-60 cell line lysates (35ug/lane). PLA2G7 (arrow) was detected using the purified Pab.



PLA2G7 Antibody (Center) (Cat. #[TA325117]) IHC analysis in formalin fixed and paraffin embedded tonsil followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the PLA2G7 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.





PLA2G7 Antibody (Center) (Cat. #[TA325117]) flow cytometric analysis of HL-60 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.