

## Product datasheet for **TA325488**

### FYN Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 1:500-1:2000
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	The antiserum was produced against A synthesized peptide derived from human Fyn
Formulation:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration:	lot specific
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific peptide.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	60 kDa
Gene Name:	FYN proto-oncogene, Src family tyrosine kinase
Database Link:	<a href="#">NP_002028</a> <a href="#">Entrez Gene 14360 Mouse</a> <a href="#">Entrez Gene 25150 Rat</a> <a href="#">Entrez Gene 2534 Human</a> <a href="#">P06241</a>

**Background:** FANCD2 Required for maintenance of chromosomal stability. Promotes accurate and efficient pairing of homologs during meiosis. Involved in the repair of DNA double-strand breaks, both by homologous recombination and single-strand annealing. May participate in S phase and G2 phase checkpoint activation upon DNA damage. Plays a role in preventing breakage and loss of missegregating chromatin at the end of cell division, particularly after replication stress. Required for the targeting, or stabilization, of BLM to non-centromeric abnormal structures induced by replicative stress.




[View online »](#)

**Synonyms:** p59-FYN; SLK; SYN

**Protein Families:** Druggable Genome, Protein Kinase

**Protein Pathways:** Adherens junction, Axon guidance, Fc epsilon RI signaling pathway, Focal adhesion, Natural killer cell mediated cytotoxicity, Pathogenic Escherichia coli infection, Prion diseases, T cell receptor signaling pathway, Viral myocarditis

**Product images:**



Western blot analysis of Fyn expression in 293 whole cell lysates. The lane on the left is treated with the antigen-specific peptide.