

## Product datasheet for TA325610

## **CD117 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** 

Recommended Dilution: WB: 1:500-1:2000 Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: lgG

Clonality: Polyclonal

Immunogen: The antiserum was produced against A synthesized peptide derived from human c-Kit

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% Formulation:

glycerol.

Concentration: lot specific

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific peptide.

Unconjugated Conjugation:

Store at -20°C as received. Storage:

Stable for 12 months from date of receipt. Stability:

**Predicted Protein Size:** 120.145 kDa

Gene Name: KIT proto-oncogene receptor tyrosine kinase

Database Link: NP 000213

Entrez Gene 16590 MouseEntrez Gene 64030 RatEntrez Gene 3815 Human

P10721

Background: KIT encodes the human homolog of the proto-oncogene c-kit. C-kit was first identified as the

> cellular homolog of the feline sarcoma viral oncogene v-kit. KIT is a type 3 transmembrane receptor for MGF (mast cell growth factor, also known as stem cell factor). Mutations in KIT are associated with gastrointestinal stromal tumors, mast cell disease, acute myelogenous

lukemia, and piebaldism.

Synonyms: C-Kit; CD117; PBT; SCFR



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Protein Families: Adult stem cells, Druggable Genome, Embryonic stem cells, ES Cell Differentiation/IPS, Protein

Kinase, Stem cell - Pluripotency, Transmembrane

**Protein Pathways:** Acute myeloid leukemia, Cytokine-cytokine receptor interaction, Endocytosis, Hematopoietic

cell lineage, Melanogenesis, Pathways in cancer

## **Product images:**



Western blot analysis of c-Kit expression in HeLa whole cell lysates, The lane on the left is treated with the antigen-specific peptide.