

## **Product datasheet for TA325618**

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

## LIM Kinase 1 (LIMK1) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** 

Recommended Dilution: WB: 1:500-1:2000 Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: lgG

Clonality: Polyclonal

Immunogen: The antiserum was produced against A synthesized peptide derived from human LIMK1

Formulation: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50%

glycerol.

Concentration: lot specific

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific peptide.

Conjugation: Unconjugated

Store at -20°C as received. Storage:

Stable for 12 months from date of receipt. Stability:

**Predicted Protein Size:** 70 kDa

Gene Name: LIM domain kinase 1

Database Link: NP 002305

Entrez Gene 16885 MouseEntrez Gene 65172 RatEntrez Gene 3984 Human

P53667





Background:


There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. LIMK1 is likely to be a component of an intracellular signaling pathway and may be involved in brain development. LIMK1 hemizygosity is implicated in the impaired visuospatial constructive cognition of Williams syndrome. Two splice variant have been identified.

Synonyms: LIMK; LIMK-1

**Protein Families:** Druggable Genome, Protein Kinase

**Protein Pathways:** Axon guidance, Fc gamma R-mediated phagocytosis, Regulation of actin cytoskeleton

## **Product images:**



Western blot analysis of LIMK1 expression in UV treated NIH-3T3 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.