

### **Product datasheet for TA325691**

#### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

## **MET Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WE

**Reactivity:** WB: 1:500-1:2000 Human, Mouse, Rat

Modifications: Phospho-specific

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** The antiserum was produced against A synthesized peptide derived from human Met around

the phosphorylation site of Tyrosine 1234

Formulation: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50%

glycerol.

**Concentration:** lot specific

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific peptide.

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 145 kDa

**Gene Name:** MET proto-oncogene, receptor tyrosine kinase

Database Link: NP 000236

Entrez Gene 17295 MouseEntrez Gene 24553 RatEntrez Gene 4233 Human

P08581

**Background:** The proto-oncogene MET product is the hepatocyte growth factor receptor and encodes

tyrosine-kinase activity. The primary single chain precursor protein is post-translationally cleaved to produce the alpha and beta subunits, which are disulfide linked to form the

mature receptor.





#### **MET Rabbit Polyclonal Antibody - TA325691**

Synonyms: AUTS9; c-Met; DFNB97; HGFR; RCCP2


**Protein Families:** Druggable Genome, Protein Kinase, Transmembrane

Protein Pathways: Adherens junction, Axon guidance, Colorectal cancer, Cytokine-cytokine receptor interaction,

Endocytosis, Epithelial cell signaling in Helicobacter pylori infection, Focal adhesion,

Melanoma, Pathways in cancer, Renal cell carcinoma

# **Product images:**



Western blot analysis of Met phosphorylation expression in HepG2 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.