

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA325692

MET Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 1:500-1:2000
Reactivity:	Human, Mouse, Rat
Modifications:	Phospho-specific
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	The antiserum was produced against A synthesized peptide derived from human Met around the phosphorylation site of Tyrosine 1349
Formulation:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration:	lot specific
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific peptide.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	145 kDa
Gene Name:	MET proto-oncogene, receptor tyrosine kinase
Database Link:	<u>NP_000236</u> <u>Entrez Gene 17295 MouseEntrez Gene 24553 RatEntrez Gene 4233 Human</u> <u>P08581</u>
Background:	The proto-oncogene MET product is the hepatocyte growth factor receptor and encodes tyrosine-kinase activity. The primary single chain precursor protein is post-translationally cleaved to produce the alpha and beta subunits, which are disulfide linked to form the mature receptor.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


	MET Rabbit Polyclonal Antibody – TA325692
--	---

Synonyms: AUTS9; c-Met; DFNB97; HGFR; RCCP2

Protein Families: Druggable Genome, Protein Kinase, Transmembrane

Protein Pathways:Adherens junction, Axon guidance, Colorectal cancer, Cytokine-cytokine receptor interaction,
Endocytosis, Epithelial cell signaling in Helicobacter pylori infection, Focal adhesion,
Melanoma, Pathways in cancer, Renal cell carcinoma

Product images:



Western blot analysis of Met phosphorylation expression in HepG2 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US