

Product datasheet for TA325751

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Glucocorticoid Receptor (NR3C1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Reactivity: WB: 1:500-1:2000

Reactivity: Human, Mouse

Modifications: Phospho-specific

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: The antiserum was produced against A synthesized peptide derived from human GR around

the phosphorylation site of Serine 203

Formulation: Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50%

glycerol.

Concentration: lot specific

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific peptide.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 86 kDa

Gene Name: nuclear receptor subfamily 3 group C member 1

Database Link: NP 000167

Entrez Gene 14815 MouseEntrez Gene 2908 Human

P04150

Background: The protein encoded by this gene is a receptor for glucocorticoids and can act as both a

transcription factor and a regulator of other transcription factors. The encoded protein can bind DNA as a homodimer or as a heterodimer with another protein such as the retinoid X receptor. This protein can also be found in heteromeric cytoplasmic complexes along with

heat shock factors and immunophilins.






Glucocorticoid Receptor (NR3C1) Rabbit Polyclonal Antibody - TA325751

Synonyms: GCCR; GCR; GCRST; GR; GRL

Protein Families: Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

Protein Pathways: Neuroactive ligand-receptor interaction

Product images:



Western blot analysis of GR phosphorylation expression in dexamethason treated A549 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.