

Product datasheet for TA325770

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PDGF Receptor beta (PDGFRB) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 1:500-1:2000

Reactivity: Human, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: The antiserum was produced against A synthesized peptide derived from human PDGFRb

Formulation: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50%

glycerol.

Concentration: lot specific

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific peptide.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 125 kDa

Gene Name: platelet derived growth factor receptor beta

Database Link: NP 002600

Entrez Gene 24629 RatEntrez Gene 5159 Human

P09619

Background: This gene encodes a cell surface tyrosine kinase receptor for members of the platelet-derived

growth factor family. These growth factors are mitogens for cells of mesenchymal origin. The

identity of the growth factor bound to a receptor monomer determines whether the functional receptor is a homodimer or a heterodimer, composed of both platelet-derived

growth factor receptor alpha and beta polypeptides.

Synonyms: CD140B; IBGC4; IMF1; JTK12; PDGFR; PDGFR-1; PDGFR1






Protein Families:

Druggable Genome, ES Cell Differentiation/IPS, Protein Kinase, Transmembrane

Protein Pathways:

Calcium signaling pathway, Colorectal cancer, Cytokine-cytokine receptor interaction, Focal adhesion, Gap junction, Glioma, MAPK signaling pathway, Melanoma, Pathways in cancer, Prostate cancer, Regulation of actin cytoskeleton

Product images:



Western blot analysis of PDGFRb expression in HepG2 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.