

## Product datasheet for **TA325807**

### NF-kB p65 (RELA) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 1:500-1:2000; IHC: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Modifications:	Phospho-specific
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	A synthesized peptide derived from human NF- $\kappa$ B p65 around the phosphorylation site of Sersine 536
Formulation:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration:	lot specific
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific peptide.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	RELA proto-oncogene, NF-kB subunit
Database Link:	<a href="#">NP_001138610</a> <a href="#">Entrez Gene 19697 Mouse</a> <a href="#">Entrez Gene 309165 Rat</a> <a href="#">Entrez Gene 5970 Human</a> <a href="#">Q04206</a>



[View online »](#)

**Background:**

Proteins encoded by the v-Rel viral oncogene and its cellular homolog, c-Rel, are members of a family of transcription factors that include the two subunits of the transcription factor NF $\kappa$ B (p50 and p65) and the Drosophila maternal morphogen, dorsal. Both proteins specifically bind to DNA sequences that are the same or slight variations of the 10 bp kB sequence in the immunoglobulin  $\gamma$  light chain enhancer. This same sequence is also present in a number of other cellular and viral enhancers. The DNA binding activity of NF $\kappa$ B is activated and rapidly transported from the cytoplasm to the nucleus in cells exposed to mitogens or growth factors. cDNAs encoding precursors for two distinct 50 kDa proteins have been described, designated p105 and p100, respectively. The p105 precursor contains p50 at its amino terminus and a C-terminal region that when expressed as a separate molecule, designated PDI, binds to p50 and regulates its activity.

**Synonyms:**


NFKB3; p65

**Protein Families:**


Druggable Genome, Transcription Factors

**Protein Pathways:**

Acute myeloid leukemia, Adipocytokine signaling pathway, Apoptosis, B cell receptor signaling pathway, Chemokine signaling pathway, Chronic myeloid leukemia, Cytosolic DNA-sensing pathway, Epithelial cell signaling in Helicobacter pylori infection, MAPK signaling pathway, Neurotrophin signaling pathway, NOD-like receptor signaling pathway, Pancreatic cancer, Pathways in cancer, Prostate cancer, RIG-I-like receptor signaling pathway, Small cell lung cancer, T cell receptor signaling pathway, Toll-like receptor signaling pathway

**Product images:**

Western blot analysis of extracts from K562 cells, using Phospho-NF- $\kappa$ B p65 (Ser536) Antibody. The lane on the left is treated with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using NF-kappaB p65 (Phospho-Ser536) Antibody.?