

## **Product datasheet for TA325957**

## p53 (TP53) Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 1:500-1:2000

Reactivity: Human

Modifications: Phospho-specific

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** The antiserum was produced against a synthesized A synthesized peptide derived from

human p53 around the phosphorylation site of Sersine 46.

Formulation: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50%

glycerol.

**Concentration:** lot specific

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific peptide.

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** tumor protein p53

Database Link: NP 000537

Entrez Gene 7157 Human

P04637



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

Tumor protein p53, a nuclear protein, plays an essential role in the regulation of cell cycle, specifically in the transition from G0 to G1. It is found in very low levels in normal cells, however, in a variety of transformed cell lines, it is expressed in high amounts, and believed to contribute to transformation and malignancy. p53 is a DNA-binding protein containing DNA-binding, oligomerization and transcription activation domains. It is postulated to bind as a tetramer to a p53-binding site and activate expression of downstream genes that inhibit growth and/or invasion, and thus function as a tumor suppressor. Mutants of p53 that frequently occur in a number of different human cancers fail to bind the consensus DNA binding site, and hence cause the loss of tumor suppressor activity. Alterations of the TP53 gene occur not only as somatic mutations in human malignancies, but also as germline mutations in some cancer-prone families with Li-Fraumeni syndrome.

**Synonyms:** BCC7; LFS1; P53; TRP53


**Protein Families:** Druggable Genome, Stem cell - Pluripotency, Transcription Factors

**Protein Pathways:** Amyotrophic lateral sclerosis (ALS), Apoptosis, Basal cell carcinoma, Bladder cancer, Cell cycle,

Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, Glioma, Huntington's disease, MAPK signaling pathway, Melanoma, Neurotrophin signaling pathway, Non-small cell lung cancer, p53 signaling pathway, Pancreatic cancer, Pathways in cancer, Prostate cancer,

Small cell lung cancer, Thyroid cancer, Wnt signaling pathway

## **Product images:**



Western blot analysis on MDA-MB-435 cell lysate using Phospho-p53 (Ser46) Antibody