

Product datasheet for TA325998

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NMDAR2B (GRIN2B) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB: 1:500-1:2000; IHC: 1:50-1:200

Reactivity: Human, Mouse, Rat **Modifications:** Phospho-specific

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: The antiserum was produced against A synthesized peptide derived from human NMDAR2B

around the phosphorylation site of Tyrosine 1336

Formulation: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50%

glycerol.

Concentration: lot specific

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific peptide.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 150 kDa

Gene Name: glutamate ionotropic receptor NMDA type subunit 2B

Database Link: NP 000825

Entrez Gene 14812 MouseEntrez Gene 24410 RatEntrez Gene 2904 Human

Q13224

Background: N-methyl-D-aspartate (NMDA) receptors are a class of ionotropic glutamate receptors. NMDA

receptor channel has been shown to be involved in long-term potentiation, an activity-dependent increase in the efficiency of synaptic transmission thought to underlie certain

kinds of memory and learning.





NMDAR2B (GRIN2B) Rabbit Polyclonal Antibody - TA325998


Synonyms: EIEE27; GluN2B; hNR3; MRD6; NMDAR2B; NR2B

Protein Families: Druggable Genome, Ion Channels: Glutamate Receptors, Transmembrane

Protein Pathways: Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Huntington's disease, Long-term

potentiation, Neuroactive ligand-receptor interaction, Systemic lupus erythematosus

Product images:



Western blot analysis of NMDAR2B phosphorylation expression in TNF treated Jurkat whole cell lysates, The lane on the left is treated with the antigen-specific peptide.