

## Product datasheet for **TA326469**

### **CACNA1G Mouse Monoclonal Antibody [Clone ID: S178A-9]**

#### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	S178A-9
Applications:	WB
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Fusion protein amino acids 2052-2172 (cytoplasmic C-terminus) of mouse Cav3.1
Formulation:	PBS pH7.4, 50% glycerol, 0.09% sodium azide
Concentration:	lot specific
Purification:	Protein G Purified
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	calcium voltage-gated channel subunit alpha1 G
Database Link:	<a href="#">NP_061496</a> <a href="#">Entrez Gene 12291 Mouse</a> <a href="#">Entrez Gene 29717 Rat</a> <a href="#">Entrez Gene 8913 Human</a> <a href="#">O43497</a>



[View online »](#)

**Background:**

Ion channels are integral membrane proteins that help establish and control the small voltage gradient across the plasma membrane of living cells by allowing the flow of ions down their electrochemical gradient. They are present in the membranes that surround all biological cells because their main function is to regulate the flow of ions across this membrane. Whereas some ion channels permit the passage of ions based on charge, others conduct based on an ionic species, such as sodium or potassium. Furthermore, in some ion channels, the passage is governed by a gate which is controlled by chemical or electrical signals, temperature, or mechanical forces. There are a few main classifications of gated ion channels. There are voltage-gated ion channels, ligand-gated, other gating systems and finally those that are classified differently, having more exotic characteristics. The first are voltage-gated ion channels which open and close in response to membrane potential. These are then separated into sodium, calcium, potassium, proton, transient receptor, and cyclic nucleotide-gated channels; each of which is responsible for a unique role. Ligand-gated ion channels are also known as ionotropic receptors, and they open in response to specific ligand molecules binding to the extracellular domain of the receptor protein. The other gated classifications include activation and inactivation by second messengers, inward-rectifier potassium channels, calcium-activated potassium channels, two-pore-domain potassium channels, light-gated channels, mechano-sensitive ion channels and cyclic nucleotide-gated channels. Finally, the other classifications are based on less normal characteristics such as two-pore channels, and transient receptor potential channels. Specifically, Calcium channel Ca<sub>v</sub>3.1 (α1G) is a low-voltage-activated T-type calcium channel. Such T-type channels are expressed throughout the body. In the heart, they may be involved in pacemaker current. In neurons, these channels may play a secondary pacemaker role. With the ubiquitous expression, it is not surprising that alterations in channel function have been implicated in disease. Drugs that act to block T-type calcium channels are used as antihypertensives, antiepileptics, and blocking of T-type calcium channels may be involved in the action of some anesthetics and antipsychotics as well. Much remains to be determined about the precise cellular localization, in vivo physiological roles, roles in disease states and possible routes to modulate their structure/function to ameliorate effects of disease.

**Synonyms:**

Ca(V)T.1; Cav3.1; NBR13

**Note:**


Detects ~200kDa. Does not cross-react with Cav3.2

**Protein Families:**

Druggable Genome, Ion Channels: Calcium, Transmembrane

**Protein Pathways:**

Calcium signaling pathway, MAPK signaling pathway, Type II diabetes mellitus

**Product images:**

Western blot analysis of Cav3.1 in rat brain membrane tissues, using a 1:1000 dilution of the antibody