

## Product datasheet for **TA326798S**

### Midkine (MDK) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ICC/IF, WB
Recommended Dilution:	WB: 1:500-1:2000
Reactivity:	Human, Mouse
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant protein of human MDK
Formulation:	Store at -20C or -80C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	midkine (neurite growth-promoting factor 2)
Database Link:	<a href="#">NP_002382</a> <a href="#">Entrez Gene 17242 Mouse</a> <a href="#">Entrez Gene 4192 Human</a> <a href="#">P21741</a>



[View online »](#)

**Background:**


Midkine, or MK, is a heparin-binding molecule involved in the regulation of growth and differentiation during embryogenesis. MK expression is tightly regulated during embryonic development by steroid receptors of the retinoic acid superfamily. The mature human MK protein is 118 amino acids in length and contains five intrachain disulfide bonds. MK is a non-glycosylated protein that shows greater than 87% identity between human and mouse. The carboxy-terminus of MK contains the principle heparin-binding site and the molecule's neurite-promoting sequences; both the amino- and carboxy-terminal sequences are required for the molecule's neurotrophic properties. An association between overexpression of MK and colon adenocarcinoma has been shown in families suffering from familial polyposis. In addition, MK functions to enhance the activity of plasminogen activator (PA).

**Synonyms:**

ARAP; MK; NEGF2

**Protein Families:**

Druggable Genome, Secreted Protein, Transmembrane

**Product images:**

Western blot analysis of extracts of SW480 cells, using MDK antibody.