

Product datasheet for TA327016S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Glycophorin C (GYPC) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, WB

Recommended Dilution: WB,1:500 - 1:2000

ELISA,Recommended starting concentration is 1 μg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Mouse, Human

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Formulation: Store at -20C or -80C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50%

glycerol, pH7.3

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 14kDa

Gene Name: glycophorin C (Gerbich blood group)

Database Link: NP 002092

Entrez Gene 71683 MouseEntrez Gene 2995 Human

P04921

Background: Glycophorin C (GYPC) is an integral membrane glycoprotein. It is a minor species carried by

human erythrocytes, but plays an important role in regulating the mechanical stability of red cells. A number of glycophorin C mutations have been described. The Gerbich and Yus phenotypes are due to deletion of exon 3 and 2, respectively. The Webb and Duch antigens, also known as glycophorin D, result from single point mutations of the glycophorin C gene. The glycophorin C protein has very little homology with glycophorins A and B. Alternate

splicing results in multiple transcript variants.






Synonyms: CD236; CD236R; GE; GPC; GPD; GYPD; PAS-2

Protein Families: Druggable Genome, Transmembrane

Product images:



Western blot analysis of various lysates using Glycophorin C (GYPC) Rabbit pAb ([TA327016]) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.