

## Product datasheet for **TA327018**

### DR5 (TNFRSF10B) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 1:500-1:2000
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant protein of human TNFRSF10B
Formulation:	PBS with 0.09% Sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	tumor necrosis factor receptor superfamily member 10b
Database Link:	<a href="#">NP_003833</a> <a href="#">Entrez Gene 8795 Human</a> <a href="#">O14763</a>



[View online »](#)

**Background:**

The tumor necrosis factor receptor family, which includes TNF-R1, Fas, DR3, DR4, DR5, and DR6, plays an important role in the regulation of apoptosis in various physiological systems. The receptors are activated by a family of cytokines that include TNF, FasL, and TRAIL. They are characterized by a highly conserved extracellular region containing cysteine-rich repeats and a conserved intracellular region of about 80 amino acids termed the death domain (DD). The DD is important for transducing the death signal by recruiting other DD containing adaptor proteins (FADD, TRADD, RIP) to the death-inducing signaling complex (DISC), resulting in activation of caspases. DR5 is a receptor for TNF-related apoptosis inducing ligand (TRAIL), which has been shown to induce apoptosis in variety of cell types and has been targeted for cancer therapy. Structurally, DR5 contains an amino-terminal leader cleavage site followed by an extracellular region containing two cysteine-rich repeats, then a central transmembrane domain and a carboxy-terminal death domain. DR5 is expressed in a wide variety of tissues and is transcriptional target for p53. It induces apoptosis through a FADD-dependent pathway. Deletion of DR5 leads to resistance in TRAIL-mediated apoptosis as well as an abrogated response to DNA-damaging stimuli.

**Synonyms:**


CD262; DR5; KILLER; TRAIL-R2; TRAILR2; TRICK2; TRICK2A; TRICK2B; TRICKB; ZTNFR9

**Protein Families:**

Druggable Genome, Transmembrane

**Protein Pathways:**

Apoptosis, Cytokine-cytokine receptor interaction, Natural killer cell mediated cytotoxicity, p53 signaling pathway

**Product images:**

Western blot analysis of HL-60 cell lysate using TNFRSF10B antibody.