

Product datasheet for **TA327037S**

Monoamine Oxidase A (MAOA) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ICC/IF, IHC, WB
Recommended Dilution:	WB 1:500 - 1:2000
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant protein of human MAOA
Formulation:	Store at -20C or -80C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	monoamine oxidase A
Database Link:	NP_000231 Entrez Gene 17161 Mouse Entrez Gene 29253 Rat Entrez Gene 4128 Human P21397



[View online »](#)

Background:

Monoamine oxidase (MAO) is an enzyme of the mitochondrial outer membrane and catalyzes the oxidative deamination of biogenic amines throughout the body. MAO is critical in the neuronal metabolism of catecholamine and indolamine transmitters. Cultured skin fibroblasts show both MAO-A and MAO-B and both MAOs differ in molecular structure. MAO-A, the primary type in fibroblasts, preferentially degrades serotonin and norepinephrine. Only MAO-B is present in platelets and only MAO-A is present in trophoblasts. MAO-B, the primary type found not only in platelets but also in the brain of man and other primates, preferentially degrades phenylethylamine and benzylamine. MAO has been of particular interest to psychiatry and genetics because of the suggestion that low activity is a genetic marker for schizophrenia. The genes which encode MAO-A and MAO-B map to human chromosome Xp11.23.

Synonyms:


BRNRS; MAO-A

Protein Families:

Druggable Genome

Protein Pathways:

Arginine and proline metabolism, Drug metabolism - cytochrome P450, Glycine, serine and threonine metabolism, Histidine metabolism, Metabolic pathways, Phenylalanine metabolism, Tryptophan metabolism, Tyrosine metabolism

Product images:

Western blot analysis of extracts of various cell lines, using MAOA antibody.