

## Product datasheet for **TA327049S**

### Factor X (F10) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ICC/IF, WB
Recommended Dilution:	WB 1:500 - 1:2000;IF 1:10 - 1:100
Reactivity:	Human, Mouse
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant protein of human F10
Formulation:	Store at -20C or -80C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	55 kDa
Gene Name:	coagulation factor X
Database Link:	<a href="#">NP_000495</a> <a href="#">Entrez Gene 14058 Mouse</a> <a href="#">Entrez Gene 2159 Human</a> <a href="#">P00742</a>



[View online »](#)

**Background:**

This gene encodes the vitamin K-dependent coagulation factor X of the blood coagulation cascade. This factor undergoes multiple processing steps before its preproprotein is converted to a mature two-chain form by the excision of the tripeptide RKR. Two chains of the factor are held together by 1 or more disulfide bonds; the light chain contains 2 EGF-like domains, while the heavy chain contains the catalytic domain which is structurally homologous to those of the other hemostatic serine proteases. The mature factor is activated by the cleavage of the activation peptide by factor IXa (in the intrinsic pathway), or by factor VIIa (in the extrinsic pathway). The activated factor then converts prothrombin to thrombin in the presence of factor Va, Ca<sup>2+</sup>, and phospholipid during blood clotting. Mutations of this gene result in factor X deficiency, a hemorrhagic condition of variable severity.

**Synonyms:**


FX; FXA

**Protein Families:**


Druggable Genome, Protease, Transmembrane

**Protein Pathways:**

Complement and coagulation cascades

**Product images:**

Western blot analysis of extracts of various cell lines, using F10 antibody.



Immunofluorescence analysis of A549 cell using F10 antibody. Blue: DAPI for nuclear staining.