

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA327054S

Peroxiredoxin 4 (PRDX4) Rabbit Polyclonal Antibody

Product data:


| Product Type: | Primary Antibodies |
|-------------------------|---|
| Applications: | ELISA, WB |
| Recommended Dilution: | WB,1:500 - 1:2000 ELISA,Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| lsotype: | lgG |
| Clonality: | Polyclonal |
| Formulation: | Store at -20C or -80C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3 |
| Concentration: | lot specific |
| Purification: | Affinity purification |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C. Avoid freeze / thaw cycles. |
| Stability: | Stable for 12 months from date of receipt. |
| Predicted Protein Size: | 31kDa |
| Gene Name: | peroxiredoxin 4 |
| Database Link: | <u>NP_006397</u> <u>Entrez Gene 53381 MouseEntrez Gene 85274 RatEntrez Gene 10549 Human</u> <u>Q13162</u> |
| Background: | The protein encoded by this gene is an antioxidant enzyme and belongs to the peroxiredoxin family. The protein is localized to the cytoplasm. Peroxidases of the peroxiredoxin family reduce hydrogen peroxide and alkyl hydroperoxides to water and alcohol with the use of reducing equivalents derived from thiol-containing donor molecules. This protein has been found to play a regulatory role in the activation of the transcription factor NF-kappaB. |
| Synonyms: | AOE37-2; AOE372; HEL-S-97n; PRX-4 |
| Protein Families: | Druggable Genome |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Product images:



Western blot analysis of various lysates using [KO Validated] Peroxiredoxin 4 (PRDX4) Rabbit pAb ([TA327054]) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Western blot analysis of lysates from wild type (WT) and Peroxiredoxin 4 (Peroxiredoxin 4 (PRDX4)) knockout (KO) 293T cells

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US