

Product datasheet for TA327068

PSMC5 Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies	
Applications:	ICC/IF, WB	
Recommended Dilution:	WB 1:500 - 1:2000;IF 1:50- 1:200	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
lsotype:	lgG	
Clonality:	Polyclonal	
Immunogen:	Recombinant protein of human PSMC5	
Formulation:	Store at -20C or -80C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3	
Concentration:	lot specific	
Purification:	Affinity purification	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Gene Name:	proteasome 26S subunit, ATPase 5	
Database Link:	<u>NP_002796</u> <u>Entrez Gene 19184 MouseEntrez Gene 81827 RatEntrez Gene 5705 Human</u> <u>P62195</u>	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

SMC5 Rabbit Polyclonal Antibody – TA327068


Background:The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure
composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4
rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are
composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6
ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase
subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration
and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An
essential function of a modified proteasome, the immunoproteasome, is the processing of
class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A
family of ATPases which have a chaperone-like activity. In addition to participation in
proteasome functions, this subunit may participate in transcriptional regulation since it has
been shown to interact with the thyroid hormone receptor and retinoid X receptor-alpha.
Two transcript variants encoding different isoforms have been found for this gene. [provided
by RefSeq, Nov 2010]

Synonyms:	p45; S8; SUG; SUG-1; SUG1; TBP10; TRIP1
Protein Families:	Druggable Genome

Proteasome

Protein Pathways:


Product images:



Western blot analysis of extracts of A431 cell and HepG2 cell lines, using PSMC5 antibody.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US





Immunofluorescence analysis of MCF7 cell using PSMC5 antibody. Blue: DAPI for nuclear staining.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US