

Product datasheet for TA327133

CDC25C Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:

Recommended Dilution: WB 1:500 - 1:2000 Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: lgG

Clonality: Polyclonal

Immunogen: Recombinant protein of human CDC25C

Store at -20C or -80C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% Formulation:

glycerol, pH7.3

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Store at -20°C as received. Storage:

Stability: Stable for 12 months from date of receipt.

Gene Name: cell division cycle 25C

Database Link: NP 001781

Entrez Gene 12532 MouseEntrez Gene 307511 RatEntrez Gene 995 Human

P30307

Background: Cdc25 is a protein phosphatase responsible for dephosphorylating and activating cdc2, a

> crucial step in regulating the entry of all eukaryotic cells into mitosis. cdc25C is constitutively phosphorylated at Ser216 throughout interphase by c-TAK1, while phosphorylation at this site is DNA damage-dependent at the G2/M checkpoint. When phosphorylated at Ser216, cdc25C binds to members of the 14-3-3 family of proteins, sequestering cdc25C in the cytoplasm and thereby preventing premature mitosis. The checkpoint kinases Chk1 and Chk2 phosphorylate

cdc25C at Ser216 in response to DNA damage.

Synonyms: CDC25: PPP1R60

Protein Families: Druggable Genome, Phosphatase, Stem cell - Pluripotency



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Pathways: Cell cycle, Oocyte meiosis, Progesterone-mediated oocyte maturation

Product images:



Western blot analysis of extracts of various cell lines, using CDC25C antibody.