

## **Product datasheet for TA327143**

## Factor XII (F12) Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** ICC/IF, WB

**Reactivity:** WB 1:500 - 1:2000 Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant protein of human F12

Formulation: Store at -20C or -80C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50%

glycerol, pH7.3

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

**Store** at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** coagulation factor XII

Database Link: NP 000496

Entrez Gene 58992 MouseEntrez Gene 306761 RatEntrez Gene 2161 Human

P00748



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


This gene encodes coagulation factor XII which circulates in blood as a zymogen. This single chain zymogen is converted to a two-chain serine protease with an heavy chain (alpha-factor XIIa) and a light chain. The heavy chain contains two fibronectin-type domains, two epidermal growth factor (EGF)-like domains, a kringle domain and a proline-rich domain, whereas the light chain contains only a catalytic domain. On activation, further cleavages takes place in the heavy chain, resulting in the production of beta-factor XIIa light chain and the alpha-factor XIIa light chain becomes beta-factor XIIa heavy chain. Prekallikrein is cleaved by factor XII to form kallikrein, which then cleaves factor XII first to alpha-factor XIIa and then to beta-factor XIIa. The active factor XIIa participates in the initiation of blood coagulation, fibrinolysis, and the generation of bradykinin and angiotensin. It activates coagulation factors VII and XI. Defects in this gene do not cause any clinical symptoms and the sole effect is that whole-blood clotting time is prolonged. [provided by RefSeq, Jul 2008]

Synonyms: HAE3; HAEX; HAF

**Protein Families:** Druggable Genome, Protease, Secreted Protein

**Protein Pathways:** Complement and coagulation cascades

## **Product images:**



Western blot analysis of extracts of Jurkat cell lines, using F12 antibody.