

Product datasheet for TA327297

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

LETMD1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, ICC/IF, IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC-P,1:50 - 1:200 IF/ICC,1:50 - 1:200

ELISA,Recommended starting concentration is 1 μg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 42kDa

Gene Name: LETM1 domain containing 1

Database Link: NP 056231

Entrez Gene 68614 MouseEntrez Gene 681352 RatEntrez Gene 25875 Human

Q6P1Q0

Background: This gene encodes a mitochondrial outer membrane protein. It has a potential role in

tumorigenesis, which may result from negative regulation of the p53 tumor suppressor gene.

Alternatively spliced transcript variants have been noted for this gene.


Synonyms: 1110019O13Rik; HCCR; HCCR-1; HCCR-2; HCCR1; HCCR2; HCRR-2

Protein Families: Druggable Genome, Transmembrane





Product images:




Western blot analysis of various lysates using LETMD1 Rabbit pAb (TA327297) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.




Immunohistochemistry analysis of paraffinembedded Human colon using LETMD1 Rabbit pAb (TA327297) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.





Immunohistochemistry analysis of paraffinembedded Mouse kidney using LETMD1 Rabbit pAb (TA327297) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffinembedded Rat testis using LETMD1 Rabbit pAb (TA327297) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.