

## **Product datasheet for TA327809**

## Paxillin (PXN) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 1:500-1:2000

Reactivity: Human, Mouse, Rat

Modifications: Phospho-specific

**Host:** Rabbit

**Isotype:** IgG

Clonality: Polyclonal

**Immunogen:** The antiserum was produced against A synthesized peptide derived from human Paxillin

around the phosphorylation site of Tyrosine 118

Formulation: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50%

glycerol.

**Concentration:** lot specific

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific peptide.

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 68 kDa

Gene Name: paxillin

Database Link: NP 002850

Entrez Gene 19303 MouseEntrez Gene 360820 RatEntrez Gene 5829 Human

P49023

**Background:** PXN is a multi-domain cytoskeletal protein that localizes primarily to focal adhesion sites to

the extracellular matrix. Phosphorylated by focal adhesion kinase (FAK) and is a component

of integrin signaling. Its phosphorylation provides docking sites for

Synonyms: FLJ16691



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




**Protein Families:** Druggable Genome, Stem cell - Pluripotency

**Protein Pathways:** Chemokine signaling pathway, Focal adhesion, Leukocyte transendothelial migration,

Regulation of actin cytoskeleton, VEGF signaling pathway

## **Product images:**



Western blot analysis of Paxillin phosphorylation expression in EGF treated HeLa whole cell lysates, The lane on the left is treated with the antigenspecific peptide.