

### **Product datasheet for TA327825**

#### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PKC delta (PRKCD) Rabbit Polyclonal Antibody

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: WE

**Recommended Dilution:** WB: 1:500-1:2000

**Reactivity:** Human, Mouse, Rat

Modifications: Phospho-specific

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: The antiserum was produced against A synthesized peptide derived from human PKCD

around the phosphorylation site of Tyrosine 64

Formulation: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50%

glycerol.

**Concentration:** lot specific

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific peptide.

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 78 kDa

**Gene Name:** protein kinase C delta

Database Link: NP 006245

Entrez Gene 18753 MouseEntrez Gene 170538 RatEntrez Gene 5580 Human

Q05655

**Background:** Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be

activated by calcium and the second messenger diacylglycerol. PKC family members

phosphorylate a wide variety of protein targets and are known to be involved

**Synonyms:** ALPS3; CVID9; MAY1; nPKC-delta; PKCD






**Protein Families:** Druggable Genome, Protein Kinase

**Protein Pathways:** Chemokine signaling pathway, Fc epsilon RI signaling pathway, Fc gamma R-mediated

phagocytosis, GnRH signaling pathway, Neurotrophin signaling pathway, Tight junction, Type

II diabetes mellitus, Vascular smooth muscle contraction

# **Product images:**



Western blot analysis of PKCD phosphorylation expression in PMA treated HepG2 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.