

Product datasheet for TA327859

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Phospholipase C gamma 1 (PLCG1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB: 1:500-1:2000; IHC: 1:50-1:200

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: The antiserum was produced against A synthesized peptide derived from human PLCG1

Formulation: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50%

glycerol.

Concentration: lot specific

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific peptide.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 150 kDa

Gene Name: phospholipase C gamma 1

Database Link: NP 002651

Entrez Gene 18803 MouseEntrez Gene 25738 RatEntrez Gene 5335 Human

P19174

Background: The protein encoded by this gene catalyzes the formation of inositol 1,4,5-trisphosphate and

diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a

cofactor and plays an important role in the intracellular transduction

Synonyms: NCKAP3; PLC-II; PLC1; PLC148; PLCgamma1

Protein Families: Druggable Genome






Protein Pathways:

Calcium signaling pathway, Epithelial cell signaling in Helicobacter pylori infection, ErbB signaling pathway, Fc epsilon RI signaling pathway, Fc gamma R-mediated phagocytosis, Glioma, Inositol phosphate metabolism, Leukocyte transendothelial migration, Metabolic pathways, Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, Nonsmall cell lung cancer, Pathways in cancer, Phosphatidylinositol signaling system, T cell receptor signaling pathway, VEGF signaling pathway, Vibrio cholerae infection

Product images:



Western blot analysis of PLCG1 expression in EGF treated COS7 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.