

## **Product datasheet for TA327913**

## **Product data:**

**Product Type:** Primary Antibodies

**CD117 Rabbit Polyclonal Antibody** 

Applications: WB

**Reactivity:** WB: 1:500-1:2000 Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** A synthesized peptide derived from human KIT.

Formulation: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50%

glycerol.Store at -20 °C.Stable for 12 months from date of receipt

**Concentration:** lot specific

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific peptide.

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 145 kDa

**Gene Name:** KIT proto-oncogene receptor tyrosine kinase

Database Link: NP 001087241

Entrez Gene 16590 MouseEntrez Gene 3815 Human

P10721

**Background:** KIT encodes the human homolog of the proto-oncogene c-kit. C-kit was first identified as the

cellular homolog of the feline sarcoma viral oncogene v-kit. KIT is a type 3 transmembrane receptor for MGF (mast cell growth factor, also known as stem cell factor). Mutations in KIT are associated with gastrointestinal stromal tumors, mast cell disease, acute myelogenous

lukemia, and piebaldism.

**Synonyms:** C-Kit; CD117; PBT; SCFR



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Protein Families: Adult stem cells, Druggable Genome, Embryonic stem cells, ES Cell Differentiation/IPS, Protein

Kinase, Stem cell - Pluripotency, Transmembrane

**Protein Pathways:** Acute myeloid leukemia, Cytokine-cytokine receptor interaction, Endocytosis, Hematopoietic

cell lineage, Melanogenesis, Pathways in cancer

## **Product images:**



Western blot analysis of KIT expression in HepG2 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.