

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA327982

## Acetylated Lysine Mouse Monoclonal Antibody [Clone ID: 15G10] Mouse Monoclonal Antibody [Clone ID: 15G10]

#### Product data:


Product Type:	Primary Antibodies
Clone Name:	15G10
Applications:	IF, WB
Recommended Dilution:	WB, IF
Reactivity:	Human
Host:	Mouse
lsotype:	IgG2b, kappa
Clonality:	Monoclonal
Immunogen:	Acetylated protein mixture
Formulation:	This antibody is provided in phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration:	lot specific
Purification:	The antibody was purified by affinity chromatography.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Background:	Proteins are reversibly and dynamically acetylated on the ε-amino group of lysine by acetyltransferases and deacetylated by deacetylases. This post-translational modification can regulate protein function (interactions with other proteins and DNA binding). Histones and transcription factors (PCAF, p53, p300, etc) appear to be the major targets of acetytransferases. Acetylation is usually associated with chromatin remodeling and transcriptional activation, although in some cases (telomeres) it is associated with gene silencing. The 15G10 antibody recognizes acetylated lysine residues on proteins.




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



### **Product images:**



Untreated Hela cells (lane 1) and sodium butyrate-treated Hela cells (24 hr treatment, lane 2) were lysed and cell extracts resolved by electrophoresis, transferred to nitrocellulose and probed with anti-acetylated lysine antibody (clone 15G10). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and a chemiluminescence system.



Untreated Hela cells (Upper Panel), or overnight nocodazole treated Hela cells (Lower Panel) stained with purified mouse monoclonal antibody against Acetylated Lysine (clone 15G10), followed by Rhodamine Red-X conjugated Donkey anti-mouse IgG and DAPI.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US