

# **Product datasheet for TA328025**

# TPX2 Mouse Monoclonal Antibody [Clone ID: 18D5]

## **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Clone Name:	18D5
Applications:	WB
Recommended Dilution:	WB, IHC, IF
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1, kappa
Clonality:	Monoclonal
Immunogen:	Amino Acid: 1-220 of human TPX2
Formulation:	This antibody is provided in phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide at 0.5 mg/ml.
Concentration:	lot specific
Purification:	The antibody was purified by affinity chromatography.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	100 kD
Gene Name:	TPX2, microtubule nucleation factor
Database Link:	<u>NP_036244</u> <u>Entrez Gene 22974 Human</u> <u>Q9ULW0</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


#### **GRIGENE** TPX2 Mouse Monoclonal Antibody [Clone ID: 18D5] – TA328025

Background:The TPX2 protein, also known as targeting protein for Xklp1, restricted expression<br/>proliferation associated protein p100, and differentially expressed in cancer and non-<br/>cancerous lung cells protein 2 (DIL2), is a 100 kD nuclear protein that contains two coiled-coil<br/>domains. The TPX2 protein is strictly associated with spindle pole and mitotic spindle during<br/>mitosis. In the G2/S position of the cell cycle, TPX2 is diffusely distributed throughout nucleus.<br/>TPX2 has been shown to be highly expressed in lung carcinomas cell lines, but not in normal<br/>lung tissues. TPX2 is thought to be required for the Ran-GTP dependent assembly of<br/>microtubules around chromosomes required to generate stable bipolar spindle with<br/>overlapping anti-parallel microtubule arrays and may also be involved in targeting Aurora-A<br/>kinase to the mitotic spindle. TPX2 has been shown to interact with a large number of<br/>proteins including serine/threonine protein kinase 6, ribosomal protein 6, Bop-1, α-tubulin,<br/>and nucleolin among others. TPX2 can be modified by phosphorylation on serine 738. The<br/>18D5 monoclonal antibody recognizes human TPX2 and has been shown to be useful for<br/>Western blotting.

 Synonyms:
 C20orf1; C20orf2; DIL-2; DIL2; FLS353; GD:C20orf1; HCA519; HCTP4; p100; REPP86

Protein Families: Druggable Genome, Stem cell - Pluripotency

### **Product images:**



MOLT-4 nuclear extracts were resolved by electrophoresis, transferred to nitrocellulose and probed with monoclonal anti-TPX2 (clone 18D5) antibody. Proteins were visualized using a goat anti-mouse secondary conjugated to HRP and a chemiluminescence detection system.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US